

Goal 3: Ensure healthy lives and promote well-being for all at all ages

Target 3.b: Support the research and development of vaccines and medicines for the communicable and non-communicable diseases that primarily affect developing countries, provide access to affordable essential medicines and vaccines, in accordance with the Doha Declaration on the TRIPS Agreement and Public Health, which affirms the right of developing countries to use to the full the provisions in the Agreement on Trade-Related Aspects of Intellectual Property Rights regarding flexibilities to protect public health, and, in particular, provide access to medicines for all

[Indicator 3.b.1: Proportion of the target population covered by all vaccines included in their national programme](#)

Institutional information

Organization(s):

World Health Organization (WHO), United Nations Children's Fund (UNICEF)

Concepts and definitions

Definition:

Coverage of DTP containing vaccine (3rd dose): Percentage of surviving infants who received the 3 doses of diphtheria and tetanus toxoid with pertussis containing vaccine in a given year.

Coverage of Measles containing vaccine (2nd dose): Percentage of children who received two dose of measles containing vaccine according to nationally recommended schedule through routine immunization services in a given year.

Coverage of Pneumococcal conjugate vaccine (last dose in the schedule): Percentage of surviving infants who received the nationally recommended doses of pneumococcal conjugate vaccine in a given year.

Coverage of HPV vaccine (last dose in the schedule): Percentage of 15 years old girls received the recommended doses of HPV vaccine.

Rationale:

This indicator aims to measure access to vaccines, including the newly available or underutilized vaccines, at the national level. In the past decades all countries added numerous new and underutilised vaccines in their national immunization schedule and there are several vaccines under final stage of development to be introduced by 2030. For monitoring diseases control and impact of vaccines it is important to measure coverage from each vaccine in national immunization schedule and the system is already in place for all national programmes, however direct measurement for proportion of population covered with all vaccines in the programme is only feasible if the country has a well-functioning national nominal immunization registry, usually an electronic one that will allow this coverage to be easily estimated. While countries will develop and strengthen immunization registries it is a need for an alternative measurement.

Concepts:

In accordance with its mandate to provide guidance to Member States on health policy matters, WHO provides global vaccine and immunization recommendations for diseases that have an international public health impact. National programmes adapt the recommendations and develop national immunization schedules, based on local disease epidemiology and national health priorities. National immunization schedules and number of recommended vaccines vary between countries, with only DTP polio and measles containing vaccines being used in all countries.

The target population for given vaccine is defined based on recommended age for administration. The primary vaccination series of most vaccines are administered in the first two years of life.

Coverage of DTP containing vaccine measure the overall system strength to deliver infant vaccination
Coverage of Measles containing vaccine ability to deliver vaccines beyond first year of life through routine immunization services.

Coverage of Pneumococcal conjugate vaccine: adaptation of new vaccines for children

Coverage of HPV vaccine: life cycle vaccination

Comments and limitations:

The rationale to select a set of vaccines reflects the ability of immunization programmes to deliver vaccines over the life cycle and to adapt new vaccines. Coverage for other WHO recommended vaccines are also available and can be provided.

Given that HPV vaccine is relatively new and vaccination schedule varies from countries to country coverage estimate will be made for girls vaccinated by age 15 and at the moment data is limited to very few countries therefore reporting will start later.

Methodology

Computation Method:

WHO and UNICEF jointly developed a methodology to estimate national immunization coverage for selected vaccines in 2000. The methodology has been refined and reviewed by expert committees over time. The methodology was published and reference is available under the reference section. Estimates time series for WHO recommended vaccines produced and published annually since 2001.

The methodology uses data reported by national authorities from countries administrative systems as well as data from immunization or multi indicator household surveys.

Disaggregation:

Geographical location, i.e. regional and national and potentially subnational estimates

Treatment of missing values:

- [At country level](#)

The first data point is the first reporting year after vaccine introduction. When country data are not available interpolation is used between 2 data points and extrapolation from the latest available data point.

- [At regional and global levels](#)

Any needed imputation is done at country level. These country values can then be used to compute regional and global ones.

Regional aggregates:

Weighted average of the country-level coverage rates where the weights are the country target population sizes based on World Population Prospects: 2017 revision from the UN Population Division. All countries from the region are included.

Sources of discrepancies:

Countries often rely on administrative coverage data, while WHO and UNICEF review and assess data from different sources including administrative systems and surveys. Differences between country produced and international estimates are mainly due to differences between coverage estimates from administrative system and survey results.

In case the vaccine is not included in national immunization schedule the coverage from private sector will not be reflected.

Data Sources

Description:

National Health Information Systems or National Immunization systems

National immunization registries

High quality household surveys with immunization module (e.g. DHS, MICS, national in-country surveys)

Collection process:

Annual data collection through established mechanism. Since 1998, in an effort to strengthen collaboration and minimize the reporting burden, WHO and UNICEF jointly collect information through a standard questionnaire (the Joint Reporting Form) sent to all Member States

http://www.who.int/immunization/monitoring_surveillance/routine/reporting/en/

Data Availability

Description:

Coverage data for different vaccines are collected annually and reviewed by WHO and UNICEF inter agency expert group and estimates made for each country and each year. Data are published both on WHO and UNICEF web sites.

http://www.who.int/immunization/monitoring_surveillance/routine/coverage/en/index4.html

<http://www.data.unicef.org/child-health/immunization>

Coverage for 2017

	DTP3	MCV2	PCV3
Global	85%	67%	44%
Australia and New Zealand	95%	92%	94%
Central Asia and Southern Asia	87%	75%	24%
Eastern Asia and South-eastern Asia	94%	90%	14%
Latin America & the Caribbean	89%	67%	77%

Northern America and Europe	93%	91%	74%
Oceania	84%	58%	75%
Sub-Saharan Africa	72%	23%	66%
Western Asia and Northern Africa (M49)	88%	84%	58%

Calendar

Data collection:

Annual data collection March-May each year. Country consultation June each year

Data release:

15 July each year for time series 1980 – release year -1. (in July 2018 estimates from 1980-2017)

Data providers

Ministries of Health, Immunization programmes

Data compilers

WHO and UNICEF

References

URL:

http://www.who.int/immunization/monitoring_surveillance/routine/coverage/en/index4.html

<https://www.unicef.org/immunization/>

References:

Burton A, Monasch R, Lautenbach B, Gacic-Dobo M, Neill M, Karimov R, Wolfson L, Jones G, Birmingham M. WHO and UNICEF estimates of national infant immunization coverage: methods and processes. Bull World Health Organ. 2009;87(7):535-41. Available at: <http://www.who.int/bulletin/volumes/87/7/08-053819/en/>

Burton A, Kowalski R, Gacic-Dobo M, Karimov R, Brown D. A Formal Representation of the WHO and UNICEF Estimates of National Immunization Coverage: A Computational Logic Approach. PLoS ONE 2012;7(10): e47806. doi:10.1371/journal.pone.0047806. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3485034/pdf/pone.0047806.pdf>

Brown D, Burton A, Gacic-Dobo M, Karimov R An Introduction to the Grade of Confidence in the WHO and UNICEF Estimates of National Immunization Coverage The Open Public Health Journal, 2013, 6, 73-76. Available at: <http://www.benthamscience.com/open/tophj/articles/V006/73TOPHJ.pdf>

Brown, David & Burton, Anthony & Gacic-Dobo, Marta. (2015). An examination of a recall bias adjustment applied to survey-based coverage estimates for multi-dose vaccines. 10.13140/RG.2.1.2086.2883.

Related indicators

Target 3.8 Achieve universal health coverage, including financial risk protection, access to quality essential health-care services and access to safe, effective, quality and affordable essential medicines and vaccines for all. Indicator 3.8.1: Coverage of essential health services (defined as the average coverage of essential services based on tracer interventions that include reproductive, maternal, newborn and child health, infectious diseases, non-communicable diseases and service capacity and access, among the general and the most disadvantaged population)